arXiv:1409.6794v1 [math.CO] 24 Sep 2014

The exterior splash in PG$(6, q)$: Transversals

S.G. Barwick and Wen-Ai Jackson
School of Mathematical Sciences, University of Adelaide
Adelaide 5005, Australia

September 25, 2014

Corresponding Author: Dr Susan Barwick, University of Adelaide, Adelaide 5005, Australia. Phone: +61 8 8313 3983, Fax: +61 8 8313 3696, email: susan.barwick@adelaide.edu.au

Keywords: Bruck-Bose representation, subplanes, exterior splash
AMS code: 51E20

Abstract

Let π be an order-q-subplane of PG$(2, q^3)$ that is exterior to ℓ_∞. Then the exterior splash of π is the set of $q^2 + q + 1$ points on ℓ_∞ that lie on an extended line of π. Exterior splashes are projectively equivalent to scattered linear sets of rank 3, covers of the circle geometry $CG(3, q)$, and hyper-reguli in PG$(5, q)$. In this article we use the Bruck-Bose representation in PG$(6, q)$ to investigate the structure of π, and the interaction between π and its exterior splash. In PG$(6, q)$, an exterior splash $\mathbb{S}$ has two sets of cover planes (which are hyper-reguli) and we show that each set has three unique transversals lines in the cubic extension PG$(6, q^3)$. These transversal lines are used to characterise the carriers of $\mathbb{S}$, and to characterise the sublines of $\mathbb{S}$.

1 Introduction

In [2, 3], the authors study order-q-subplanes of PG$(2, q^3)$ and determine their representation in the Bruck-Bose representation in PG$(6, q)$. A full characterisation in PG$(6, q)$ was given for order-q-subplanes that are secant or tangent to ℓ_∞ in PG$(2, q^3)$. However, an order-q-subplane of PG$(2, q^3)$ that is exterior to ℓ_∞ forms a complex structure in PG$(6, q)$, and one of our original motivations was to investigate this structure in PG$(6, q)$. As part

of this investigation, we studied the splash of an order-q-subplane, and found this to be a structure rich in detail, and related to many other structures in geometry, see [4, 5, 6, 18]. In this article we study the structure of an exterior order-q-subplane in $\mathrm{PG}(6,q)$, as well as the interplay between this structure and the associated exterior splash.

Let π be a subplane of $\mathrm{PG}(2,q^3)$ of order q that is exterior to ℓ_∞. The lines of π meet ℓ_∞ in a set $\mathbb{S}$ of size q^2+q+1, called the *exterior splash* of π. Properties of the exterior splash are studied in [6]. In particular, we showed that the sets of points in an exterior splash has arisen in many different situations, namely scattered linear sets of rank 3, covers of the circle geometry $CG(3,q)$, hyper-reguli in $\mathrm{PG}(5,q)$, and Sherk surfaces of size q^2+q+1.

This article proceeds as follows. In Section 2 we introduce the notation we use for the Bruck-Bose representation of $\mathrm{PG}(2,q^3)$ in $\mathrm{PG}(6,q)$, as well as presenting some other preliminary results. In Section 3 we choose an order-q-subplane $\mathcal{B}$ in $\mathrm{PG}(2,q^3)$ that is exterior to ℓ_∞, and calculate its points and exterior splash. This order-q-subplane will be used in many of the proofs in this article. In Section 4, we study the structure of an order-q-subplane in $\mathrm{PG}(6,q)$. In particular, we show that it contains q^2+q+1 twisted cubics; it is the intersection of nine quadrics; and has an interesting tangent plane at each point.

We next study the exterior splash $\mathbb{S}$ of ℓ_∞ in the Bruck-Bose representation in $\mathrm{PG}(5,q)$. By results of Bruck [10], $\mathbb{S}$ has two switching sets denoted $\mathbb{C},\mathbb{T}$, which we call covers of $\mathbb{S}$. The three sets $\mathbb{S},\mathbb{C},\mathbb{T}$ are called hyper-reguli in [19]. In Section 5, we find coordinates for the sets $\mathbb{S},\mathbb{C},\mathbb{T}$. In Section 6, we show that each of the sets $\mathbb{S},\mathbb{C},\mathbb{T}$ has a unique triple of conjugate transversal lines in the cubic extension $\mathrm{PG}(5,q^3)$. These nine transversal lines are used to characterise the carriers of an exterior splash. Further, these transversal lines are related to the set of $q-1$ disjoint splashes of ℓ_∞ that have common carriers, and their related covers. We interpret this result in terms of replacing hyper-reguli to create André planes. In Section 7 we use the transversal lines to characterise the order-q-sublines of an exterior splash in terms of how the corresponding 2-reguli meet the cover planes.

2 The Bruck-Bose representation

2.1 The Bruck-Bose representation of $\mathrm{PG}(2,q^3)$ in $\mathrm{PG}(6,q)$

We introduce the notation we will use for the Bruck-Bose representation of $\mathrm{PG}(2,q^3)$ in $\mathrm{PG}(6,q)$. We work with the finite field $\mathbb{F}_q=\mathrm{GF}(q)$, and let $\mathbb{F}_q'=\mathbb{F}_q\backslash\{0\}$.

A 2-*spread* of $\mathrm{PG}(5,q)$ is a set of q^3+1 planes that partition $\mathrm{PG}(5,q)$. A 2-*regulus* of $\mathrm{PG}(5,q)$ is a set of $q+1$ mutually disjoint planes $\pi_1,\ldots,\pi_{q+1}$ with the property that if a line meets three of the planes, then it meets all $q+1$ of them. A 2-regulus $\mathcal{R}$ has a set of q^2+q+1 mutually disjoint *ruling lines* that meet every plane of $\mathcal{R}$. A 2-regulus

is uniquely determined by three mutually disjoint planes, or four (ruling) lines (mutually disjoint and lying in general position). A 2-spread $\mathcal{S}$ is *regular* if for any three planes in $\mathcal{S}$, the 2-regulus containing them is contained in $\mathcal{S}$. See [16] for more information on 2-spreads.

The following construction of a regular 2-spread of $\mathrm{PG}(5,q)$ will be needed. Embed $\mathrm{PG}(5,q)$ in $\mathrm{PG}(5,q^3)$ and let g be a line of $\mathrm{PG}(5,q^3)$ disjoint from $\mathrm{PG}(5,q)$. Let g^q, g^{q^2} be the conjugate lines of g; both of these are disjoint from $\mathrm{PG}(5,q)$. Let P_i be a point on g; then the plane $\langle P_i, P_i^q, P_i^{q^2}\rangle$ meets $\mathrm{PG}(5,q)$ in a plane. As P_i ranges over all the points of g, we get q^3+1 planes of $\mathrm{PG}(5,q)$ that partition $\mathrm{PG}(5,q)$. These planes form a regular 2-spread $\mathcal{S}$ of $\mathrm{PG}(5,q)$. The lines g, g^q, g^{q^2} are called the (conjugate skew) *transversal lines* of the 2-spread $\mathcal{S}$. Conversely, given a regular 2-spread in $\mathrm{PG}(5,q)$, there is a unique set of three (conjugate skew) transversal lines in $\mathrm{PG}(5,q^3)$ that generate $\mathcal{S}$ in this way.

We will use the linear representation of a finite translation plane $\mathcal{P}$ of dimension at most three over its kernel, due independently to André [1] and Bruck and Bose [11, 12]. Let Σ_∞ be a hyperplane of $\mathrm{PG}(6,q)$ and let $\mathcal{S}$ be a 2-spread of Σ_∞. We use the phrase *a subspace of* $\mathrm{PG}(6,q)\backslash\Sigma_\infty$ to mean a subspace of $\mathrm{PG}(6,q)$ that is not contained in Σ_∞. Consider the following incidence structure: the *points* of $\mathcal{A}(\mathcal{S})$ are the points of $\mathrm{PG}(6,q)\backslash\Sigma_\infty$; the *lines* of $\mathcal{A}(\mathcal{S})$ are the 3-spaces of $\mathrm{PG}(6,q)\backslash\Sigma_\infty$ that contain an element of $\mathcal{S}$; and *incidence* in $\mathcal{A}(\mathcal{S})$ is induced by incidence in $\mathrm{PG}(6,q)$. Then the incidence structure $\mathcal{A}(\mathcal{S})$ is an affine plane of order q^3. We can complete $\mathcal{A}(\mathcal{S})$ to a projective plane $\mathcal{P}(\mathcal{S})$; the points on the line at infinity ℓ_∞ have a natural correspondence to the elements of the 2-spread $\mathcal{S}$. The projective plane $\mathcal{P}(\mathcal{S})$ is the Desarguesian plane $\mathrm{PG}(2,q^3)$ if and only if $\mathcal{S}$ is a regular 2-spread of $\Sigma_\infty \cong \mathrm{PG}(5,q)$ (see [8]). For the remainder of this article, we use $\mathcal{S}$ to denote a regular 2-spread of $\Sigma_\infty \cong \mathrm{PG}(5,q)$.

We use the following notation. If T is a point of ℓ_∞ in $\mathrm{PG}(2,q^3)$, we use $[T]$ to refer to the plane of $\mathcal{S}$ corresponding to T. More generally, if X is a set of points of $\mathrm{PG}(2,q^3)$, then we let $[X]$ denote the corresponding set in $\mathrm{PG}(6,q)$. If P is an affine point of $\mathrm{PG}(2,q^3)$, we generally simplify the notation and also use P to refer to the corresponding affine point in $\mathrm{PG}(6,q)$, although in some cases, to avoid confusion, we use $[P]$.

When $\mathcal{S}$ is a regular 2-spread, we can relate the coordinates of $\mathcal{P}(\mathcal{S}) \cong \mathrm{PG}(2,q^3)$ and $\mathrm{PG}(6,q)$ as follows. Let τ be a primitive element in $\mathbb{F}_{q^3}$ with primitive polynomial $x^3 - t_2x^2 - t_1x - t_0$. Every element $\alpha \in \mathbb{F}_{q^3}$ can be uniquely written as $\alpha = a_0 + a_1\tau + a_2\tau^2$ with $a_0, a_1, a_2 \in \mathbb{F}_q$. Points in $\mathrm{PG}(2,q^3)$ have homogeneous coordinates (x,y,z) with $x,y,z \in \mathbb{F}_{q^3}$, not all zero. Let the line at infinity ℓ_∞ have equation $z=0$; so the affine points of $\mathrm{PG}(2,q^3)$ have coordinates $(x,y,1)$. Points in $\mathrm{PG}(6,q)$ have homogeneous coordinates $(x_0,x_1,x_2,y_0,y_1,y_2,z)$ with $x_0,x_1,x_2,y_0,y_1,y_2,z \in \mathbb{F}_q$. Let Σ_∞ have equation $z=0$. Let $P=(\alpha,\beta,1)$ be a point of $\mathrm{PG}(2,q^3)$. We can write $\alpha = a_0 + a_1\tau + a_2\tau^2$ and $\beta = b_0 + b_1\tau + b_2\tau^2$ with $a_0,a_1,a_2,b_0,b_1,b_2 \in \mathbb{F}_q$. We want to map the element α of $\mathbb{F}_{q^3}$ to the vector (a_0,a_1,a_2), and we use the following notation to do this:

$$[\alpha] = (a_0, a_1, a_2).$$

This gives us some notation for the Bruck-Bose map, denoted ϵ, from an affine point $P = (\alpha, \beta, 1) \in \mathrm{PG}(2, q^3) \backslash \ell_\infty$ to the corresponding affine point $[P] \in \mathrm{PG}(6, q) \backslash \Sigma_\infty$, namely

$$\epsilon(\alpha, \beta, 1) = [(\alpha, \beta, 1)] = ([\alpha], [\beta], 1) = (a_0, a_1, a_2, b_0, b_1, b_2, 1).$$

More generally, if $z \in \mathbb{F}_q$, then $\epsilon(\alpha, \beta, z) = ([\alpha], [\beta], z) = (a_0, a_1, a_2, b_0, b_1, b_2, z)$.

Consider the case when $z = 0$, a point on ℓ_∞ in $\mathrm{PG}(2, q^3)$ has coordinates $L = (\alpha, \beta, 0)$ for some $\alpha, \beta \in \mathbb{F}_{q^3}$. In $\mathrm{PG}(6, q)$, the point $\epsilon(\alpha, \beta, 0) = ([\alpha], [\beta], 0)$ is one point in the spread element $[L]$ corresponding to L. Moreover, the spread element $[L]$ consists of all the points $\{([\alpha x], [\beta x], 0) : x \in \mathbb{F}'_{q^3}\}$. Hence the regular 2-spread $\mathcal{S}$ consists of the planes $\{[kx], [x], 0] : x \in \mathbb{F}'_{q^3}\}$ for $k \in \mathbb{F}_{q^3} \cup \{\infty\}$.

With this coordinatisation for the Bruck-Bose map, we can calculate the coordinates of the transversal lines of the regular 2-spread $\mathcal{S}$.

Lemma 2.1 [2] *Let* $p_0 = t_1 + t_2\tau - \tau^2 = -\tau^q\tau^{q^2}$, $p_1 = t_2 - \tau = \tau^q + \tau^{q^2}$, $p_2 = -1$, *and* $A = (p_0, p_1, p_2)$. *Then in the cubic extension* $\mathrm{PG}(6, q^3)$, *one transversal line of the regular* 2*-spread* $\mathcal{S}$ *contains the two points* $A_1 = (p_0, p_1, p_2, 0, 0, 0, 0) = (A, [0], 0)$ *and* $A_2 = (0, 0, 0, p_0, p_1, p_2, 0) = ([0], A, 0)$.

2.2 Some useful homographies

In order to simplify the notation in some of the following coordinate based proofs, we define some homographies which will be useful. We can represent an element $x = x_0 + x_1\tau + x_2\tau^2 \in \mathbb{F}_{q^3}$ as a point $[x] = (x_0, x_1, x_2)$ in $\mathrm{PG}(2, q)$. For $k \in \mathbb{F}'_{q^3}$, consider the homography ζ_k in $\mathrm{PGL}(3, q)$ with matrix M_k that maps $[x]$ to $[kx]$. Let $k \in \mathbb{F}'_{q^3}$, write $k = k_0 + k_1\tau + k_2\tau^2$, then $M_k = k_0M_1 + k_1M_\tau + k_2M_{\tau^2}$, and hence

$$M_kA = kA \qquad \text{and} \qquad M_kA^{q^2} = k^{q^2}A^{q^2}, \tag{1}$$

where $A = (p_0, p_1, p_2)^t$ is defined in Lemma 2.1. We use ζ_k to define the homography θ_k of $\mathrm{PG}(5, q)$, $k \in \mathbb{F}_{q^3}$:

$$\theta_k \colon ([x], [y]) \to ([kx], [y]) = (M_k[x], [y]).$$

From the matrix M_τ, we construct three more homographies of $\mathrm{PG}(2, q)$ with matrices U_0, U_1, U_2 that help with the notation in the proof of Theorem 7.3. For $i = 0, 1, 2$, (with p_i as in Lemma 2.1), let

$$U_i = (p_0I + p_1M_\tau + p_2M_\tau^2)^{q^i} = \begin{pmatrix} p_0^{q^i} & \tau^{q^i}p_0^{q^i} & \tau^{2q^i}p_0^{q^i} \\ p_1^{q^i} & \tau^{q^i}p_1^{q^i} & \tau^{2q^i}p_1^{q^i} \\ p_2^{q^i} & \tau^{q^i}p_2^{q^i} & \tau^{2q^i}p_2^{q^i} \end{pmatrix}.$$

Then

$$U_i \begin{pmatrix} a_0 \\ a_1 \\ a_2 \end{pmatrix} = (a_0 + a_1\tau^{q^i} + a_2\tau^{2q^i}) \begin{pmatrix} p_0^{q^i} \\ p_1^{q^i} \\ p_2^{q^i} \end{pmatrix}, a_0, a_1, a_2 \in \mathbb{F}_{q^3}.$$

Note that if $a_0, a_1, a_2 \in \mathbb{F}_q$, and $\alpha = a_0 + a_1\tau + a_2\tau^2$, then $[\alpha] = (a_0, a_1, a_2)^t$, and we write the matrix equation as $U_i[\alpha] = \alpha^{q^i} A^{q^i}$.

2.3 Sublines in the Bruck-Bose representation

An *order-q-subplane* of $\mathrm{PG}(2, q^3)$ is a subplane of $\mathrm{PG}(2, q^3)$ of order q. Equivalently, it is an image of $\mathrm{PG}(2, q)$ under $\mathrm{PGL}(3, q^3)$. An *order-q-subline* of $\mathrm{PG}(2, q^3)$ is a line of an order-q-subplane of $\mathrm{PG}(2, q^3)$. An *order-q-subline* of $\mathrm{PG}(1, q^3)$ is defined to be one of the images of $\mathrm{PG}(1, q) = \{(a, 1) : a \in \mathbb{F}_q\} \cup \{(1, 0)\}$ under $\mathrm{PGL}(2, q^3)$.

In [2, 3], the authors determine the representation of order-q-subplanes and order-q-sublines of $\mathrm{PG}(2, q^3)$ in the Bruck-Bose representation in $\mathrm{PG}(6, q)$, and we quote the results for order-q-sublines which are needed in this article. We first introduce some terminology to simplify the statements. Recall that $\mathcal{S}$ is a regular 2-spread in the hyperplane at infinity Σ_∞ in $\mathrm{PG}(6, q)$.

Definition 2.2 *1. An* $\mathcal{S}$-special conic *is a non-degenerate conic $\mathcal{C}$ contained in a plane of $\mathcal{S}$, such that the extension of $\mathcal{C}$ to $\mathrm{PG}(6, q^3)$ meets the transversals of $\mathcal{S}$.*

2. An $\mathcal{S}$-special twisted cubic *is a twisted cubic $\mathcal{N}$ in a 3-space of $\mathrm{PG}(6, q)\backslash\Sigma_\infty$ about a plane of $\mathcal{S}$, such that the extension of $\mathcal{N}$ to $\mathrm{PG}(6, q^3)$ meets the transversals of $\mathcal{S}$.*

Theorem 2.3 [2] *Let b be an order-q-subline of $\mathrm{PG}(2, q^3)$.*

1. *If $b \subset \ell_\infty$, then in $\mathrm{PG}(6, q)$, b corresponds to a 2-regulus of $\mathcal{S}$. Conversely every 2-regulus of $\mathcal{S}$ corresponds to an order-q-subline of ℓ_∞.*

2. *If b meets ℓ_∞ in a point, then in $\mathrm{PG}(6, q)$, b corresponds to a line of $\mathrm{PG}(6, q)\backslash\Sigma_\infty$. Conversely every line of $\mathrm{PG}(6, q)\backslash\Sigma_\infty$ corresponds to an order-q-subline of $\mathrm{PG}(2, q^3)$ tangent to ℓ_∞.*

3. *If b is disjoint from ℓ_∞, then in $\mathrm{PG}(6, q)$, b corresponds to an $\mathcal{S}$-special twisted cubic. Further, a twisted cubic $\mathcal{N}$ of $\mathrm{PG}(6, q)$ corresponds to an order-q-subline of $\mathrm{PG}(2, q^3)$ if and only if $\mathcal{N}$ is $\mathcal{S}$-special.*

The article [2] also determines the representation of secant and tangent order-q-subplanes of $\mathrm{PG}(2, q^3)$ in $\mathrm{PG}(6, q)$. The representation of an exterior order-q-subplane in $\mathrm{PG}(6, q)$

is more complex to describe. One of the motivations of this work was to investigate this representation in more detail. Some aspects of the representation are discussed in more detail in Section 4.

2.4 Properties of exterior splashes

We need the following group theoretic results about order-q-subplanes and exterior splashes which appear in [6].

Theorem 2.4 *Consider the collineation group $G = \mathrm{PGL}(3, q^3)$ acting on $\mathrm{PG}(2, q^3)$. The subgroup G_ℓ fixing a line ℓ is transitive on the order-q-subplanes that are exterior to ℓ, and is transitive on the exterior splashes of ℓ.*

Theorem 2.5 *The group $K = \mathrm{PGL}(3, q^3)_\pi$ acting on $\mathrm{PG}(2, q^3)$ and fixing an order-q-subplane π is transitive on the points of π.*

Consider the group $G = \mathrm{PGL}(3, q^3)$ acting on $\mathrm{PG}(2, q^3)$. It has an important subgroup $I = G_{\pi,\ell}$ which fixes an order-q-subplane π, and a line ℓ exterior to π. By [6], I fixes exactly three lines: ℓ, and its conjugates m, n with respect to π; and I fixes exactly three points: $E_1 = \ell \cap m$, $E_2 = \ell \cap n$, $E_3 = m \cap n$, which are conjugate with respect to π. This group I identifies two special points $E_1 = \ell \cap m$, $E_2 = \ell \cap n$ on ℓ which are called the *carriers* of the exterior splash $\mathbb{S}$ of π. This is consistent with the definition of carriers of a circle geometry $CG(3, q)$, see [6]. Further, the fixed points and fixed lines of $I = G_{\pi,\ell}$ are used to define an important class of conics in an order-q-subplane π with respect to an exterior line ℓ.

Definition 2.6 *Let π be an order-q-subplane with exterior line ℓ, and let $I = \mathrm{PGL}(3, q^3)_{\pi,\ell}$ have fixed lines ℓ, m, n and fixed points E_1, E_2, E_3.*

1. *A conic of π whose extension to $\mathrm{PG}(2, q^3)$ contains the three points E_1, E_2, E_3 is called a* (π, ℓ)-special conic *of π.*
2. *A dual conic of π whose extension to $\mathrm{PG}(2, q^3)$ contains the three lines ℓ, m, n is called a* (π, ℓ)-special dual conic.

Note that special conics and special dual conics are necessarily irreducible (see [3]). This definition of (π, ℓ)-special conics is consistent with the definition of $\mathcal{S}$-special conics given in Definition 2.2. That is, if $\mathcal{C}$ is an $\mathcal{S}$-special conic of a spread element $\alpha \in \mathcal{S}$, and $P = \alpha \cap g$, where g is a transversal of the regular 2-spread $\mathcal{S}$, then $\mathcal{C}$ is an (α, PP^q)-special conic of α.

3 Coordinatising an exterior order-q-subplane

Recall from Theorem 2.4 that the group of homographies of $\mathrm{PG}(2,q^3)$ is transitive on pairs (π,ℓ) where π is an order-q-subplane exterior to the line ℓ. So if we want to use coordinates to prove a result about exterior order-q-subplanes, we can without loss of generality prove it for a particular exterior order-q-subplane. In this section we calculate the coordinates for an order-q-subplane $\mathcal{B}$ of $\mathrm{PG}(2,q^3)$ that is exterior to ℓ_∞. We will use this order-q-subplane to analyse the exterior splash in the Bruck-Bose representation in $\mathrm{PG}(6,q)$.

We begin with the order-q-subplane $\pi_0 = \mathrm{PG}(2,q)$ of $\mathrm{PG}(2,q^3)$. Recall that ℓ_∞ has equation $z=0$, so π_0 is secant to ℓ_∞. We give a homography that maps π_0 to an order-q-subplane exterior to ℓ_∞. We define the following matrices:

$$K = \begin{pmatrix} -\tau & 1 & 0 \\ -\tau^q & 1 & 0 \\ \tau\tau^q & -\tau-\tau^q & 1 \end{pmatrix}, \qquad K' = \begin{pmatrix} -1 & 1 & 0 \\ -\tau^q & \tau & 0 \\ -\tau^{2q} & \tau^2 & \tau-\tau^q \end{pmatrix}. \tag{2}$$

Let σ be the homography of $\mathrm{PG}(2,q^3)$ with matrix K. Note that as KK' is a $\mathbb{F}_{q^3}$-multiple of the identity matrix, it follows that K' is a matrix for the inverse homography σ^{-1}. Thus, if we write the points X of $\mathrm{PG}(2,q^3)$ as column vectors, and the lines ℓ of $\mathrm{PG}(2,q^3)$ as row vectors, then $\sigma\colon X \mapsto KX$ and $\sigma\colon \ell \mapsto \ell K'$.

Theorem 3.1 *In $\mathrm{PG}(2,q^3)$, let $\pi_0 = \mathrm{PG}(2,q)$, and let $\mathcal{B} = \sigma(\pi_0)$, then*

1. *$\mathcal{B}$ is an order-q-subplane exterior to ℓ_∞.*
2. *The exterior splash of $\mathcal{B}$ on ℓ_∞ is*
$$\mathbb{S} = \{(k,1,0) : k \in \mathbb{F}_{q^3}, k^{q^2+q+1}=1\} \equiv \{(\tau^{(q-1)i},1,0) : 0 \le i < q^2+q+1\}.$$
3. *The carriers of the exterior splash of $\mathcal{B}$ are $E_1=(1,0,0)$ and $E_2=(0,1,0)$.*

Proof Note that σ maps $\pi_0 = \mathrm{PG}(2,q)$ to $\mathcal{B}$ and maps the line $\ell = [-\tau\tau^q, \tau+\tau^q, -1]$ to $\ell_\infty = [0,0,1]$. By [6, Lemma 2.4], π_0 is exterior to ℓ and has carriers $E=(1,\tau,\tau^2)$ and $E^q = (1,\tau^q,\tau^{2q})$ on ℓ. Hence $\mathcal{B}$ is exterior to ℓ_∞ and has carriers $\sigma(E) = (0,1,0)$ and $\sigma(E^q) = (1,0,0)$ on ℓ_∞.

By considering the action of σ on the lines $[l,m,n]$ ($l,m,n \in \mathbb{F}_q$, not all zero) of π_0, we calculate the lines of $\mathcal{B}$ are $\ell_{l,m,n} = [-l-\tau^q m-\tau^{2q}n,\ l+\tau m+\tau^2 n,\ n(\tau-\tau^q)]$, $l,m,n \in \mathbb{F}_q$, not all zero. The exterior splash of $\mathcal{B}$ consists of the points $Q_{l,m,n} = \ell_{l,m,n} \cap \ell_\infty = (l+\tau m+\tau^2 n,\ (l+\tau m+\tau^2 n)^q, 0)$. Writing $y = l+\tau m+\tau^2 n$, gives $Q_{l,m,n} \equiv (y,y^q,0) \equiv (y^{1-q},1,0)$ and writing $y=\tau^{-j}$ for some $j \in \{0,\dots,q^3-2\}$ yields $Q_{l,m,n} \equiv (\tau^{j(q-1)},1,0)$. Note that if we write $j = n(q^2+q+1)+i$ where $0 \le i < q^2+q+1$, then $\tau^{j(q-1)} = \tau^{i(q-1)}$. So we may assume that $Q_{l,m,n} = (\tau^{i(q-1)},1,0)$ with $0 \le i < q^2+q+1$. Finally, note that the solutions to $k^{q^2+q+1}=1$ are $\tau^{i(q-1)}$, $0 \le i < q^2+q+1$. □

4 The structure of the subplane in $\mathrm{PG}(6,q)$

If π is an exterior order-q-subplane of $\mathrm{PG}(2,q^3)$, then in the Bruck-Bose representation in $\mathrm{PG}(6,q)$, π corresponds to a set of q^2+q+1 affine points denoted $[\pi]$. It is difficult to characterise the structure of $[\pi]$. We note that as π contains q^2+q+1 order-q-sublines that are exterior to ℓ_∞, then by Theorem 2.3, $[\pi]$ contains q^2+q+1 $\mathcal{S}$-special twisted cubics, each lying in a 3-space through a distinct plane of the exterior splash of π. In this section we aim to determine more about the structure of $[\pi]$.

4.1 An exterior splash is the intersection of nine quadrics

We show that the structure $[\pi]$ of $\mathrm{PG}(6,q)$ corresponding to an exterior order-q-subplane π of $\mathrm{PG}(2,q^3)$ is the intersection of nine quadrics in $\mathrm{PG}(6,q)$. This is analogous to [4, Theorem 9.2] which shows that a *tangent* order-q-subplane of $\mathrm{PG}(2,q^3)$ corresponds to a structure in $\mathrm{PG}(6,q)$ that is the intersection of nine quadrics.

Theorem 4.1 *Let π be an exterior order-q-subplane in $\mathrm{PG}(2,q^3)$. Then in $\mathrm{PG}(6,q)$, the corresponding set $[\pi]$ is the intersection of nine quadrics.*

Proof By Theorem 2.4, we can without loss of generality prove this for the order-q-subplane $\mathcal{B}$ coordinatised in Section 3. We use the homographies σ, σ^{-1} with matrices K, K' respectively, given in (2). A point $P=(x,y,1)\in\mathrm{PG}(2,q^3)$ belongs to $\mathcal{B}$ if its pre-image $K'P=(-x+y,\ -\tau^q x+\tau y,\ -\tau^{2q}x+\tau^2 y+(\tau-\tau^q))$ belongs to $\pi_0=\mathrm{PG}(2,q)$. Suppose firstly that $-x+y\neq 0$, then

$$K'P\equiv\left(1,\ \frac{-\tau^q x+\tau y}{-x+y},\ \frac{-\tau^{2q}x+\tau^2 y+(\tau-\tau^q)}{-x+y}\right).$$

This belongs to $\pi_0=\mathrm{PG}(2,q)$ if and only if the second and third coordinates belong to $\mathbb{F}_q$, that is,

$$\left(\frac{-\tau^q x+\tau y}{-x+y}\right)^q = \frac{-\tau^q x+\tau y}{-x+y}, \tag{3}$$

$$\left(\frac{-\tau^{2q}x+\tau^2 y+(\tau-\tau^q)}{-x+y}\right)^q = \frac{-\tau^{2q}x+\tau^2 y+(\tau-\tau^q)}{-x+y}. \tag{4}$$

Writing $x=x_0+x_1\tau+x_2\tau^2$ and $y=y_0+y_1\tau+y_2\tau^2$, where $x_i,y_i\in\mathbb{F}_q$, $i=1,2,3$, then equating powers of $1,\tau,\tau^2$, yields three quadratic equations from each condition, a total of six, each of which represents a quadric in $\mathrm{PG}(6,q)$.

Secondly, suppose $-\tau^q x+\tau y\neq 0$, then

$$K'P\equiv\left(\frac{-x+y}{-\tau^q x+\tau y},\ 1,\ \frac{-\tau^{2q}x+\tau^2 y+(\tau-\tau^q)}{-\tau^q x+\tau y}\right).$$

As before, this lies in π_0 if and only if

$$\left(\frac{-x+y}{-\tau^q x+\tau y}\right)^q = \frac{-x+y}{-\tau^q x+\tau y}, \quad (5)$$

$$\left(\frac{-\tau^{2q}x+\tau^2 y+(\tau-\tau^q)}{-\tau^q x+\tau y}\right)^q = \frac{-\tau^{2q}x+\tau^2 y+(\tau-\tau^q)}{-\tau^q x+\tau y}, \quad (6)$$

leading to a further six quadrics in $\mathrm{PG}(6,q)$. The equations (3) and (5) give the same triple of quadrics. Hence the point P lies in $\mathcal{B}$ if and only if the point $[P]$ lies on a total of nine quadrics in $\mathrm{PG}(6,q)$. Finally, note that if both $-x+y=0$ and $-\tau^q x+\tau y=0$, then $x=y=0$ and the point P has coordinates $(0,0,1)$. This satisfies all the quadratic equations from (3), (4), (6), and so in $\mathrm{PG}(6,q)$, $[P]$ lies on each of the nine quadrics. $\square$

4.2 Tangent planes at points of an exterior subplane

We now consider a point P lying in an exterior order-q-subplane π of $\mathrm{PG}(2,q^3)$. In the Bruck-Bose representation in $\mathrm{PG}(6,q)$, P corresponds to an affine point which we also denote by P. We show that in $\mathrm{PG}(6,q)$, there is a unique *tangent plane* $\mathcal{T}_P$ at P to the structure $[\pi]$. We show that there are two equivalent ways to define this tangent plane. Recall from Theorem 2.3 that the order-q-sublines of π correspond to twisted cubics in $\mathrm{PG}(6,q)$. Theorem 4.2 shows that we can define $\mathcal{T}_P$ by looking at the tangent lines at P to these twisted cubics. Then Theorem 4.3 shows that we can define $\mathcal{T}_P$ by looking at the tangent space of P with respect to the nine quadrics defined by $[\pi]$.

Theorem 4.2 *Let π be an exterior order-q-subplane of $\mathrm{PG}(2,q^3)$, and let P be a point of π. Label the lines of π through P by $\ell_0,\ldots,\ell_q$. In $\mathrm{PG}(6,q)$, ℓ_i corresponds to a twisted cubic $[\ell_i]$. Let m_i be the unique tangent line to $[\ell_i]$ through P. Then the lines $m_0,\ldots,m_q$ lie in a plane $\mathcal{T}_P$, called the tangent plane of $[\pi]$ at P.*

Proof By Theorems 2.4 and 2.5, we can without loss of generality prove this for the order-q-subplane $\mathcal{B}$ coordinatised in Section 3, and the point $P=(0,0,1)$ of $\mathcal{B}$. First consider the order-q-subplane $\pi_0=\mathrm{PG}(2,q)$. The point $P=(0,0,1)$ lies in π_0, and the lines of π_0 through P have coordinates $\ell'_m=[m,1,0]$, $m\in\mathbb{F}_q\cup\{\infty\}$. Points on the line ℓ'_m distinct from P have coordinates $P'_x=(1,-m,x)$ for $x\in\mathbb{F}_q$. We map the plane π_0 to $\mathcal{B}$ using the homography σ with matrix K given in (2). As $\sigma(P)=P$, the lines of $\mathcal{B}$ through P are $\ell_m=\sigma(\ell'_m)$, $m\in\mathbb{F}_q\cup\{\infty\}$. Points on the line ℓ_m distinct from P have coordinates $P_x=\sigma(P'_x)=(-\tau-m,\ -\tau^q-m,\ \tau\tau^q+(\tau+\tau^q)m+x)$, for $x\in\mathbb{F}_q$.

To convert this to a coordinate in $\mathrm{PG}(6,q)$, we need to multiply by an element of $\mathbb{F}_{q^3}$ so that the last coordinate lies in $\mathbb{F}_q$. Let $F(x)=\tau\tau^q+(\tau+\tau^q)m+x$ (the third coordinate in P_x). As $F(x)\in\mathbb{F}_{q^3}$, we have $F(x)^{q^2+q+1}\in\mathbb{F}_q$. So in $\mathrm{PG}(6,q)$, we have the point $P_x=([-(\tau+m)F(x)^{q^2+q}],\ [-(\tau^q+m)F(x)^{q^2+q}],\ F(x)^{q^2+q+1})$.

By Theorem 2.3, the line ℓ_m of $\mathrm{PG}(2,q^3)$ corresponds to a twisted cubic $[\ell_m] = \{P_x : x \in \mathbb{F}_q\} \cup \{P\}$ of $\mathrm{PG}(6,q)$. Consider the unique tangent to $[\ell_m]$ through P, and let I_m be the intersection of this tangent with Σ_∞. We will show that the points I_m, $m \in \mathbb{F}_q \cup \{\infty\}$ form a line. To find the point of intersection I_m, we let $Q_x = PP_x \cap \Sigma_\infty$, then let $x \to \infty$ and calculate $I_m = Q_\infty$.

$$\begin{aligned} I_m &= \lim_{x\to\infty} PP_x \cap \Sigma_\infty = \lim_{x\to\infty} ([-(\tau+m)F(x)^{q^2+q}], [-(\tau^q+m)F(x)^{q^2+q}], 0) \\ &= ([-(\tau+m)],\ -[\tau^q+m], 0). \end{aligned}$$

Hence the points I_m, $m \in \mathbb{F}_q \cup \{\infty\}$ form a line $\ell = \langle([1],[1],0),([\tau],[\tau^q],0)\rangle$ in Σ_∞. Hence the tangent lines $m_0, \ldots, m_q$ to the twisted cubics of $[\pi]$ through P form a plane $\mathcal{T}_P = \langle \ell, P\rangle$ through P, as required. □

Theorem 4.3 *Let π be an exterior order-q-subplane of $\mathrm{PG}(2,q^3)$, and let P be a point of π. In $\mathrm{PG}(6,q)$, consider the intersection of the tangent spaces at P of the nine quadrics corresponding to $[\pi]$. Then this intersection is equal to the tangent plane $\mathcal{T}_P$ of $[\pi]$ at P as defined in Theorem 4.2.*

Proof By Theorems 2.4 and 2.5, we can without loss of generality prove this for the order-q-subplane $\mathcal{B}$ coordinatised in Section 3, and the point $P = (0,0,1)$ of $\mathcal{B}$. In $\mathrm{PG}(6,q)$, consider the nine quadrics corresponding to $[\mathcal{B}]$ which are given in equations (4), (5) and (6). We want to find the set of lines through P that meet each of these nine quadrics twice at P. Every line ℓ of $\mathrm{PG}(6,q)$ through P has form $\ell = RP$ for some point $R = ([u],[v],0) \in \Sigma_\infty$, $u, v \in \mathbb{F}_{q^3}$. So the points of ℓ are of the form $P_s = P + sR = ([su],[sv],1)$ where $s \in \mathbb{F}_q$. Substituting the point P_s into the quadrics of (4) gives

$$(-\tau^{2q}su + \tau^{2q}sv + (\tau - \tau^q))^q(-su+sv) \;=\; (-\tau^{2q}su + \tau^2 sv + (\tau-\tau^q))(-su+sv)^q.$$

This expression is a polynomial of degree two in s. The line $\ell = PR$ is tangent to the three quadrics of (4) if this expression has a repeated root $s = 0$, that is, if the coefficient of s is equal to zero. That is, $(\tau-\tau^q)^q(-u+v) = (\tau-\tau^q)(-u+v)^q$, and so $k = (-u+v)/(\tau-\tau^q)$ is in $\mathbb{F}_q$. Rearranging gives $v = k(\tau-\tau^q)+u$. Substituting the point P_s into the quadrics of (5) gives no constraints. Substituting the point P_s into the quadrics of (6) and simplifying gives the constraint $m = (-\tau^q u + \tau v)/(\tau-\tau^q)$ lies in $\mathbb{F}_q$, and so $v = (m(\tau-\tau^q)+\tau^q u)/\tau$. Equating this with the expression for v obtained from (4) above gives $u = m - k\tau$, and so $v = m - k\tau^q$. Hence the line $\ell = PR$ is tangent to all nine quadrics when R has form

$$R = ([u],[v],0) = ([m-k\tau],[m-k\tau^q],0) = m([1],[1],0) - k([\tau],[\tau^q],0).$$

Thus the tangent space to $[\mathcal{B}]$ at P is the plane through P and the line $\ell = \langle([1],[1],0),([\tau],[\tau^q],0)\rangle$ of Σ_∞. This is the same as the tangent plane $\mathcal{T}_P$ to $[\mathcal{B}]$ at P calculated in the proof of Theorem 4.2. □

5 Coordinatising the exterior splash and its covers

Let $\mathbb{S}$ be an exterior splash of $\mathrm{PG}(1,q^3)$. In the Bruck-Bose representation, $\mathbb{S}$ corresponds to a set of q^2+q+1 planes of the regular 2-spread $\mathcal{S}$ in $\Sigma_\infty \cong \mathrm{PG}(5,q)$. To simplify the notation, we use the same symbol $\mathbb{S}$ to denote both the points of the exterior splash on ℓ_∞, and the planes of the exterior splash contained in $\mathcal{S}$. In [6], we show that an exterior splash is projectively equivalent to a cover of the circle geometry $\mathrm{CG}(3,q)$. Hence by Bruck [10], there are two *switching sets* $\mathbb{X}$, $\mathbb{Y}$ for $\mathbb{S}$. That is, $\mathbb{X}$ and $\mathbb{Y}$ consist of q^2+q+1 planes each, such that the planes of the three sets $\mathbb{S}$, $\mathbb{X}$ and $\mathbb{Y}$ each cover the same set of points. Further, planes from different sets meet in unique points, and planes in the same set are disjoint. The three sets $\mathbb{S},\mathbb{X},\mathbb{Y}$ are called *hyper-reguli* in [13, 19]. In this article, we call the families $\mathbb{X}$ and $\mathbb{Y}$ *covers of the exterior splash* $\mathbb{S}$.

In this section we take the order-q-subplane $\mathcal{B}$ coordinatised in Section 3, with exterior splash $\mathbb{S}$, and use [19] to calculate the coordinates of the two covers of $\mathbb{S}$. We denote the covers corresponding to $\mathcal{B}$ by $\mathbb{T}$ and $\mathbb{C}$. The cover denoted $\mathbb{T}$ in the following lemma is called the *tangent cover* of $\mathbb{S}$ with respect to $\mathcal{B}$, or the tangent cover of $[\mathcal{B}]$. The cover denoted $\mathbb{C}$ in the following lemma is called *conic cover* of $\mathbb{S}$ with respect to $\mathcal{B}$, or the conic cover of $[\mathcal{B}]$. The nomenclature for tangent covers comes from Theorem 5.3 which shows that the tangent planes $\mathcal{T}_P$ of $[\mathcal{B}]$ meet Σ_∞ in lines that lie in distinct planes of the cover $\mathbb{T}$. The nomenclature for the conic cover comes from [7] which shows that the planes in the cover $\mathbb{C}$ are related to the $(\mathcal{B},\ell_\infty)$-special conics in $\mathcal{B}$.

Lemma 5.1 *Let $\mathbb{S}$ be the exterior splash of the exterior order-q-subplane $\mathcal{B}$ coordinatised in Section 3. Let $\mathcal{K}=\{k=\tau^{i(q-1)} : 0\le i<q^2+q+1\}$. In $\mathrm{PG}(6,q)$, $\mathbb{S}$ and its two covers $\mathbb{T},\mathbb{C}$ have planes given by*

$$\begin{aligned}
\mathbb{S} &= \{[S_k]=\{([kx],[x],0) : x\in\mathbb{F}'_{q^3}\} : k\in\mathcal{K}\}\\
\mathbb{T} &= \{[T_k]=\{([kx],[x^q],0) : x\in\mathbb{F}'_{q^3}\} : k\in\mathcal{K}\}\\
\mathbb{C} &= \{[C_k]=\{([kx],[x^{q^2}],0) : x\in\mathbb{F}'_{q^3}\} : k\in\mathcal{K}\}.
\end{aligned}$$

Proof The points of ℓ_∞ in $\mathrm{PG}(2,q^3)$ have coordinates $S_k=(k,1,0)$ for $k\in\mathbb{F}_{q^3}\cup\{\infty\}$. Hence in the Bruck-Bose representation of ℓ_∞ in $\Sigma_\infty\cong\mathrm{PG}(5,q)$, planes of the regular 2-spread $\mathcal{S}$ are given by $[S_k]=\{([kx],[x]) : x\in\mathbb{F}'_{q^3}\}$, for $k\in\mathbb{F}_{q^3}\cup\{\infty\}$. Consider the homography β (of order 3) of $\Sigma_\infty\cong\mathrm{PG}(5,q)$ defined by:

$$\beta : ([x],[y])\to([x],[y^q]). \tag{7}$$

We consider the action of β on the planes of $[S_k]$. For each $k\in\mathbb{F}_{q^3}\cup\{\infty\}$, define the planes $[T_k],[C_k]$ by $\beta([S_k])=[T_k]$ and $\beta([T_k])=[C_k]$. That is, $[T_k]=\{([kx],[x^q]) : x\in\mathbb{F}'_{q^3}\}$, and $[C_k]=\{([kx],[x^{q^2}]) : x\in\mathbb{F}'_{q^3}\}$.

We now consider the exterior order-q-subplane $\mathcal{B}$ coordinatised in Section 3 which by Theorem 3.1 has exterior splash $\mathbb{S} = \{S_k = (k,1,0) : k \in \mathcal{K}\} \subset \ell_\infty$, and carriers $S_\infty = (1,0,0)$, $S_0 = (0,1,0)$. Note that in $\mathrm{PG}(5,q)$, the carriers of $\mathcal{B}$ lie in each of the three sets of planes, as $[S_0] = [T_0] = [C_0]$ and $[S_\infty] = [T_\infty] = [C_\infty]$. In $\mathrm{PG}(5,q)$, we have $\mathbb{S} = \{[S_k] : k \in \mathcal{K}\}$. Let $\mathbb{T} = \{[T_k] : k \in \mathcal{K}\}$ and $\mathbb{C} = \{[C_k] : k \in \mathcal{K}\}$, then $\beta : \mathbb{S} \mapsto \mathbb{T} \mapsto \mathbb{C}$. By [19], the sets $\mathbb{S}$, $\mathbb{T}$, $\mathbb{C}$ cover the same set of points. Moreover, planes in the same set are disjoint, and planes from different sets meet in one point. That is, $\mathbb{T}$ and $\mathbb{C}$ are the two covers of $\mathbb{S}$. □

The next lemma demonstrates a useful homography of $\mathrm{PG}(6,q)$ that acts regularly on the cover planes in each cover of $\mathcal{B}$. Recall that τ is a zero of the primitive polynomial $x^3 - t_2x^2 - t_1x - t_0$.

Lemma 5.2 *Let $\mathbb{S}$ be the exterior splash of the exterior order-q-subplane $\mathcal{B}$ coordinatised in Section 3 with covers $\mathbb{C}$ and $\mathbb{T}$ coordinatised in Lemma 5.1. Consider the homography $\Theta \in \mathrm{PGL}(7,q)$ with 7×7 matrix*

$$\begin{pmatrix} M & 0 & 0 \\ 0 & M & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{where} \quad M = \begin{pmatrix} 0 & 0 & t_0 \\ 1 & 0 & t_1 \\ 0 & 1 & t_2 \end{pmatrix}.$$

Then in $\mathrm{PG}(6,q)$, Θ fixes each plane of the regular 2-spread $\mathcal{S}$; maps the cover plane $[C_k] \in \mathbb{C}$ to $[C_{\tau^{1-q}k}] \in \mathbb{C}$; and maps the cover plane $[T_k] \in \mathbb{T}$ to $[T_{\tau^{1-q^2}k}] \in \mathbb{T}$, $k \in \mathcal{K}$.

Proof It is straightforward to show that Θ fixes the planes of the regular 2-spread $\mathcal{S}$ (so it also fixes the planes of the exterior splash $\mathbb{S}$). In fact $\langle\Theta\rangle$ acts regularly on the set of points, and on the set of lines, of each spread element. Note that M is the matrix M_τ defined in Section 2.2, and so $M[x] = [\tau x]$. Consider the action of Θ on a point of the cover plane $[C_k] \in \mathbb{C}$ coordinatised in Lemma 5.1. We have $([kx],[x^{q^2}],0)^\Theta = ([\tau kx],[\tau x^{q^2}],0) \equiv ([\tau^{1-q}k(\tau^q x)],[(\tau^q x)^{q^2}],0)$ which lies in the cover plane $[C_{\tau^{1-q}k}]$ of $\mathbb{C}$. Similarly a point $([kx],[x^q],0)$ in the cover plane $[T_k] \in \mathbb{T}$ maps under Θ to the point $([\tau^{1-q^2}k(\tau^{q^2}x)],[(\tau^{q^2}x)^q],0)$ which lies in the cover plane $[T_{\tau^{1-q^2}k}]$ of $\mathbb{T}$. □

Theorem 5.3 *Let P be a point of an exterior order-q-subplane π. In $\mathrm{PG}(6,q)$, the tangent plane $\mathcal{T}_P$ at P to $[\pi]$ meets Σ_∞ in a line that lies in a plane of the tangent cover $\mathbb{T}$ of $[\pi]$. Moreover, distinct points of π correspond to distinct cover planes of $\mathbb{T}$.*

Proof By Theorems 2.4 and 2.5, we can without loss of generality prove this result for the order-q-subplane $\mathcal{B}$ coordinatised in Section 3 and the point $P = (0,0,1) \in \mathcal{B}$. In $\mathrm{PG}(6,q)$, let $\mathcal{T}_P$ be the tangent plane at P, the line $\ell = \mathcal{T}_P \cap \Sigma_\infty$ was calculated in the proof of Theorem 4.2 to be $\ell = \{a([1],[1],0) + b([\tau],[\tau^q],0) : a,b \in \mathbb{F}_q\}$. The points of ℓ all lie in the plane $[T_1] = \{[x],[x^q],0) | x \in \mathbb{F}'_{q^3}\}$, which by Theorem 5.1 is a plane of the

tangent cover $\mathbb{T}$ of $\mathcal{B}$. The collineation of Lemma 5.2 is transitive on the cover planes of $\mathbb{T}$, hence each cover plane contains a line of a distinct tangent plane. Hence there is a one-to-one correspondence between points of π and cover planes of $\mathbb{T}$. □

6 Transversal lines of covers

Recall that a regular 2-spread in $\mathrm{PG}(5,q)$ has three (conjugate skew) transversals in $\mathrm{PG}(5,q^3)$ which meet each (extended) plane of $\mathcal{S}$. In this section we consider an exterior splash $\mathbb{S} \subset \mathcal{S}$, and show in Lemma 6.1 that the transversals of the 2-spread $\mathcal{S}$ are the only lines of $\mathrm{PG}(5,q^3)$ that meet every extended plane of $\mathbb{S}$. We then consider the two sets of cover planes $\mathbb{T}$ and $\mathbb{C}$. We show in Theorem 6.2 that each can be extended to regular 2-spread, and we calculate the coordinates of the corresponding transversal lines in Theorem 6.3. We then show in Theorem 6.5 that the nine transversals of $\mathbb{S}$, $\mathbb{C}$ and $\mathbb{T}$ can be used to characterise the carriers of the exterior splash $\mathbb{S}$. Finally, in Theorem 6.6, we discuss the transversal lines in the situation when ℓ_∞ is partitioned into exterior splashes with common carriers.

6.1 The exterior splash and its covers have unique transversals

If $\mathcal{X}$ is a set in $\mathrm{PG}(6,q)$ (such as a line, a plane, or a conic), then we denote its natural extension to $\mathrm{PG}(6,q^3)$ by $\mathcal{X}^*$. Let $\mathcal{S}$ be the regular 2-spread in Σ_∞ of the Bruck-Bose representation in $\mathrm{PG}(6,q)$. If we extend the planes of $\mathcal{S}$ to $\mathrm{PG}(6,q^3)$, yielding $\mathcal{S}^*$, then there are exactly three transversal lines to $\mathcal{S}^*$, that is, three lines that meet every plane of $\mathcal{S}^*$. These three lines are conjugate and skew. We now consider an exterior splash $\mathbb{S} \subset \mathcal{S}$ and extend the planes of $\mathbb{S}$ to $\mathrm{PG}(6,q^3)$, yielding $\mathbb{S}^*$. We show that there are exactly three lines of $\mathrm{PG}(6,q^3)$ that meet every plane of $\mathbb{S}^*$, namely the three transversals of $\mathcal{S}$.

Lemma 6.1 *Let $\mathcal{S}$ be a regular 2-spread in $\mathrm{PG}(5,q)$, and let $\mathbb{S} \subset \mathcal{S}$ be an exterior splash. In the cubic extension $\mathrm{PG}(5,q^3)$, there are exactly three transversals to $\mathbb{S}$, namely the three transversals of $\mathcal{S}$. Hence $\mathbb{S}$ lies in a unique regular 2-spread, namely $\mathcal{S}$.*

Proof The three conjugate transversal lines of the regular 2-spread $\mathcal{S}$, denoted $g_{\mathbb{S}}, g_{\mathbb{S}}^q, g_{\mathbb{S}}^{q^2}$, are also transversals of $\mathbb{S}$. Suppose there is a fourth transversal line ℓ of $\mathbb{S}$. Then the four lines $g_{\mathbb{S}}, g_{\mathbb{S}}^q, g_{\mathbb{S}}^{q^2}, \ell$ are pairwise skew. Further, these four lines are ruling lines of a unique 2-regulus $\mathcal{R}$ of $\Sigma_\infty^* \cong \mathrm{PG}(5,q^3)$, which contains the set of extended planes $\mathbb{S}^*$. Now consider two planes $[L],[M] \in \mathbb{S}$, the corresponding points L, M of ℓ_∞ in $\mathrm{PG}(2,q^3)$ lie in two order-q-sublines contained in $\mathbb{S}$ by [17, Corollary 15]. Hence by Theorem 2.3, $[L],[M]$ lie in two 2-reguli $\mathcal{R}_1, \mathcal{R}_2$ which are contained in $\mathbb{S}$. Let P be a point in $[L]$, then there are unique lines m_1, m_2 through P that are ruling lines of $\mathcal{R}_1, \mathcal{R}_2$ respectively. Now $\mathcal{R}_1$, $\mathcal{R}_2$

lie in $\mathbb{S}$, and so lie in $\mathcal{R}$, so the extended lines m_i^*, $i = 1, 2$, are two ruling lines of $\mathcal{R}$ that meet in a point P, a contradiction. Hence the line ℓ cannot exist. That is, there are only three transversal lines to $\mathbb{S}$, and these are necessarily the transversals of $\mathcal{S}$. □

Analogous to Lemma 6.1 about an exterior splash $\mathbb{S}$, we will show that the two covers of $\mathbb{S}$ each have exactly three conjugate transversal lines in $\mathrm{PG}(6, q^3)$.

Theorem 6.2 *In* $\mathrm{PG}(5, q)$*, let* $\mathbb{S}$ *be an exterior splash with covers* $\mathbb{T}$ *and* $\mathbb{C}$*. Then in the cubic extension* $\mathrm{PG}(5, q^3)$*,*

1. *The cover* $\mathbb{T}$ *has exactly three transversal lines:* $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2} \subseteq \mathrm{PG}(5, q^3) \backslash \mathrm{PG}(5, q)$*, and so* $\mathbb{T}$ *lies in a unique regular* 2*-spread.*

2. *The cover* $\mathbb{C}$ *has exactly three transversal lines:* $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2} \subseteq \mathrm{PG}(5, q^3) \backslash \mathrm{PG}(5, q)$*, and so* $\mathbb{C}$ *lies in a unique regular* 2*-spread.*

Proof By Theorem 2.4, we can without loss of generality prove this for the exterior splash of the order-q-subplane $\mathcal{B}$ coordinatised in Section 3. We use the coordinatisation in $\mathrm{PG}(5, q)$ of the exterior splash $\mathbb{S}$ of $\mathcal{B}$ and the two covers $\mathbb{T}$, $\mathbb{C}$ given in Lemma 5.1. The homography β of $\mathrm{PG}(5, q)$ defined in (7) can be regarded as a homography of $\mathrm{PG}(5, q^3)$. By the proof of Lemma 5.1, β maps the spread plane $[S_k]$, $k \in \mathbb{F}_{q^3} \cup \{\infty\}$ to a plane $[T_k]$ of Σ_∞ (note that $[T_k] \in \mathbb{T}$ if $k \in \mathcal{K}$). Thus β maps the transversal line $g_{\mathbb{S}}$ to a line $g_{\mathbb{T}}$ that meets every extended plane $[T_k]^*$, $k \in \mathbb{F}_{q^3} \cup \{\infty\}$. Hence $g_{\mathbb{T}}$ meets each (extended) cover plane of $\mathbb{T} = \{[T_k] : k \in \mathcal{K}\}$, and so $g_{\mathbb{T}}$ is a transversal line of $\mathbb{T}$. Thus the conjugate lines $g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$ are also transversals of $\mathbb{T}$. The argument that these are the only transversals of $\mathbb{T}$ is similar to the proof of Lemma 6.1. Finally, note that the set $\{[T_k] : k \in \mathbb{F}_{q^3} \cup \{\infty\}\}$ is a regular 2-spread with transversals $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$, and so it is the unique regular 2-spread containing $\mathbb{T}$. As β maps the plane $[T_k]$ to the plane $[C_k]$, and maps the line $g_{\mathbb{T}}$ to the line $g_{\mathbb{C}}$, a similar result holds for the conic cover $\mathbb{C}$. □

Later we will need the coordinates of the point of intersection of the transversal lines with the corresponding cover planes, and we calculate these next.

Theorem 6.3 *Let* $\mathcal{B}$ *be the order-*q*-subplane coordinatised in Section* 3 *with exterior splash* $\mathbb{S}$ *and covers* $\mathbb{C}, \mathbb{T}$*. Let* $p_0 = t_1 + t_2\tau - \tau^2 = -\tau^q\tau^{q^2}$*,* $p_1 = t_2 - \tau = \tau^q + \tau^{q^2}$*,* $p_2 = -1$*, and* $\eta = p_0 + p_1\tau + p_2\tau^2$*. Let* $A_1 = (p_0, p_1, p_2, 0, 0, 0, 0)$*,* $A_2 = (0, 0, 0, p_0, p_1, p_2, 0)$*. Then in* $\mathrm{PG}(6, q^3)$*,*

1. *One transversal line of* $\mathbb{S}$ *is* $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$*, and* $g_{\mathbb{S}} \cap [S_k]^* = kA_1 + A_2$*.*

2. *One transversal line of* $\mathbb{T}$ *is* $g_{\mathbb{T}} = \langle A_1, A_2^{q^2} \rangle$*, and* $g_{\mathbb{T}} \cap [T_k]^* = kA_1 + \eta^{1-q^2} A_2^{q^2}$*.*

3. One transversal line of $\mathbb{C}$ *is* $g_{\mathbb{C}} = \langle A_1, A_2^q \rangle$*, and* $g_{\mathbb{C}} \cap [C_k]^* = kA_1 + \eta^{1-q} A_2^q$*.*

Proof We use the coordinatisation in $\mathrm{PG}(5,q)$ of the exterior splash $\mathbb{S}$ of $\mathcal{B}$ and the two covers $\mathbb{T}$, $\mathbb{C}$ given in Lemma 5.1. Lemma 2.1 shows that $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$ is a transversal line for the regular 2-spread $\mathcal{S}$, where $A_1 = (p_0, p_1, p_2, 0, 0, 0) = (A, [0])$ and $A_2 = (0, 0, 0, p_0, p_1, p_2) = ([0], A)$. Hence $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$ is a transversal line for the exterior splash $\mathbb{S}$. The planes of the regular 2-spread $\mathcal{S}$ are $[S_k] = \{([kx], [x]) : x \in \mathbb{F}'_{q^3}\}$, $k \in \mathbb{F}_{q^3} \cup \{\infty\}$. We first show that the extended plane $[S_k]^*$ meets the line $g_{\mathbb{S}}$ in the point $kA_1 + A_2$. Consider the point $P = p_0([k],[1]) + p_1([k\tau],[\tau]) + p_2([k\tau^2],[\tau^2])$ of $\mathrm{PG}(5,q^3)$, and note that $P \in [S_k]^*$. Using the matrix M_k defined in Section 2.2, we have $P = p_0(M_k[1],[1]) + p_1(M_k[\tau],[\tau]) + p_2(M_k[\tau^2],[\tau^2]) = (M_k A, A) = (kA, A)$ by (1). Hence $P = kA_1 + A_2$ which lies in $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$, that is, P is the intersection of $g_{\mathbb{S}}$ and $[S_k]^*$ proving part 1.

Consider the homography β defined in (7), acting on $\mathrm{PG}(5,q^3)$. The proof of Theorem 6.2 shows that β maps $g_{\mathbb{S}}$ to $g_{\mathbb{T}}$, and maps $g_{\mathbb{T}}$ to $g_{\mathbb{C}}$. Each element $y \in \mathbb{F}'_{q^3}$ can be considered as a point $[y]$ in $\mathrm{PG}(2,q)$. The collineation of $\mathrm{PG}(2,q)$ mapping the point $[y]$ to $[y^q]$ is a homography, and can be represented using a matrix N with entries in $\mathbb{F}_q$. We omit the transpose notation, and write $N[y] = [y^q]$. Hence we can write the collineation β as $\beta([x],[y]) = ([x], N[y])$. Clearly $\beta(A_1) = A_1$, we show that $\beta(A_2) = A_2^{q^2}$. Recall the point $A = (p_0, p_1, p_2) = p_0[1] + p_1[\tau] + p_2[\tau^2]$, so $NA = p_0[1] + p_1[\tau^q] + p_2[\tau^{2q}]$. Using the matrix M_k from Section 2.2, it is straightforward to write this as $NA = \left(p_0^{q^2} I + p_1^{q^2} M_\tau + p_2^{q^2} M_{\tau^2}\right)^q [1]$. Now $(p_0^{q^2} I + p_1^{q^2} M_\tau + p_2^{q^2} M_{\tau^2})[1] = A^{q^2}$, and $(p_0^{q^2} I + p_1^{q^2} M_\tau + p_2^{q^2} M_{\tau^2}) A^{q^2} = \eta^{q^2} A^{q^2}$ by (1). So repeated use of (1) yields $NA = \eta^{q^2(q-1)} A^{q^2} = \eta^{1-q^2} A^{q^2}$. Further, as N is over $\mathbb{F}_q$, we have

$$NA = \eta^{1-q^2} A^{q^2}, \quad NA^q = \eta^{q-1} A, \quad NA^{q^2} = \eta^{q^2-q} A^q. \tag{8}$$

Hence $\beta(kA_1 + A_2) = kA_1 + \eta^{1-q^2} A_2^{q^2}$. As $\beta : g_{\mathbb{S}} \mapsto g_{\mathbb{T}}$, we have $g_{\mathbb{T}} \cap [T_k]^* = kA_1 + \eta^{1-q^2} A_2^{q^2}$ and $g_{\mathbb{T}} = \langle A_1, A_2^{q^2} \rangle$, proving part 2. Similarly, calculating $\beta(kA_1 + \eta^{1-q^2} A_2^{q^2}) = kA_1 + \eta^{1-q^2+q^2-q} A_2^q = kA_1 + \eta^{1-q} A_2^q$, and using $\beta : g_{\mathbb{T}} \mapsto g_{\mathbb{C}}$, we get $g_{\mathbb{C}} \cap [C_k]^* = kA_1 + \eta^{1-q} A_2^q$ and $g_{\mathbb{C}} = \langle A_1, A_2^q \rangle$. □

We can use the transversals of the covers $\mathbb{T}$ and $\mathbb{C}$ to generalise the notion of $\mathcal{S}$-special conics and twisted cubics in $\mathrm{PG}(6,q)$ defined in Definition 2.2. We define $\mathbb{C}$-special here, $\mathbb{T}$-special is similarly defined.

Definition 6.4 *1. A* $\mathbb{C}$-special conic *is a non-degenerate conic* $\mathcal{C}$ *contained in a plane of* $\mathbb{C}$*, such that the extension of* $\mathcal{C}$ *to* $\mathrm{PG}(6,q^3)$ *meets the transversals of* $\mathbb{C}$*.*

2. A $\mathbb{C}$-special twisted cubic *is a twisted cubic* $\mathcal{N}$ *in a 3-space of* $\mathrm{PG}(6,q) \backslash \Sigma_\infty$ *about a plane of* $\mathbb{C}$*, such that the extension of* $\mathcal{N}$ *to* $\mathrm{PG}(6,q^3)$ *meets the transversals of* $\mathbb{C}$*.*

6.2 Characterising the carriers of an exterior splash in $\mathrm{PG}(6,q)$

Let $\mathcal{S}$ be a regular 2-spread of $\mathrm{PG}(5,q)$, and let $\mathbb{S}$ be an exterior splash contained in $\mathcal{S}$, with covers $\mathbb{C}$ and $\mathbb{T}$. Then we can characterise the carriers of $\mathbb{S}$ in terms of the nine transversals of $\mathbb{S}$, $\mathbb{C}$ and $\mathbb{T}$.

Theorem 6.5 *Let $\mathcal{S}$ be a regular 2-spread of $\mathrm{PG}(5,q)$, and let $\mathbb{S} \subset \mathcal{S}$ be an exterior splash with covers $\mathbb{C}$, $\mathbb{T}$, whose corresponding triples of transversal lines are $g_{\mathbb{S}}, g_{\mathbb{S}}^q, g_{\mathbb{S}}^{q^2}$, $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$, $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$ respectively. Then the carriers of $\mathbb{S}$ are the only two planes of $\mathcal{S}$ whose extension to $\mathrm{PG}(5,q^3)$ meets all nine transversal lines.*

Proof By Theorem 2.4, we can without loss of generality show this for the exterior splash $\mathbb{S}$ of the exterior order-q-subplane $\mathcal{B}$ coordinatised in Section 3, with carriers $E_1 = (1,0,0)$, $E_2 = (0,1,0)$. In $\mathrm{PG}(6,q)$, the transversal lines $g_{\mathbb{S}}, g_{\mathbb{S}}^q, g_{\mathbb{S}}^{q^2}$ each meet the carriers $[E_1]$, $[E_2]$ of $\mathbb{S}$. We use the notation for planes $[S_k] \in \mathcal{S}$, $[T_k] \in \mathbb{T}$ and $[C_k] \in \mathbb{C}$ from Lemma 5.1. By the proof of Theorem 6.2, in the cubic extension $\mathrm{PG}(5,q^3)$, the transversal lines $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$ meet each plane $[T_k]$, $k \in \mathbb{F}_{q^3} \cup \{\infty\}$; and the transversal lines $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$ meet each plane $[C_k]$, $k \in \mathbb{F}_{q^3} \cup \{\infty\}$. The carriers of $\mathbb{S}$ satisfy $[E_2] = [S_0] = [T_0] = [C_0]$ and $[E_1] = [S_\infty] = [T_\infty] = [C_\infty]$. Hence in the cubic extension $\mathrm{PG}(5,q^3)$, all nine transversal lines meet the carriers of $\mathbb{S}$.

We now show that no other plane of the regular 2-spread $\mathcal{S}$ meets all nine transversal lines. We use the homography with matrix M_k defined in Section 2.2. A plane of the regular 2-spread $\mathcal{S}$ distinct from $[E_1]$, $[E_2]$ has form $[S_k] = \{([kx],[x],0) : x \in \mathbb{F}'_{q^3}\}$, for some $k \in \mathbb{F}'_{q^3}$. This plane is spanned by the three points $S_{0,k} = ([k],[1],0) = (M_k(1,0,0),(1,0,0))$, $S_{1,k} = ([k\tau],[\tau],0) = (M_k(0,1,0),(0,1,0))$ and $S_{2,k} = ([k\tau^2],[\tau^2],0) = (M_k(0,0,1),(0,0,1))$. Hence the extension $[S_k]^*$ to $\mathrm{PG}(5,q^3)$ contains the points $S_{k,j} = c_0S_{0,j} + c_1S_{1,j} + c_2S_{2,j}$ where $c_i \in \mathbb{F}_{q^3}$, not all zero. By Theorem 6.3, a general point X on the transversal line $g_{\mathbb{T}}$ has coordinates $X = rA_1 + A_2^{q^2} = (rp_0, rp_1, rp_2, p_0^{q^2}, p_1^{q^2}, p_2^{q^2})$, for some $r \in \mathbb{F}_{q^3} \cup \{\infty\}$. Now $S_{j,k} = X$ if and only if $c_i = p_i^{q^2}$, $i = 0,1,2$, and $M_k(c_0,c_1,c_2) = r(p_0,p_1,p_2)$. That is, $M_kA^{q^2} = rA$. However, by (1), $M_kA^{q^2} = k^{q^2}A^{q^2}$, so there are no solutions to c_0, c_1, c_2. Hence the transversal line $g_{\mathbb{T}}$ does not meet any further plane of the regular 2-spread $\mathcal{S}$, and so $g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$ do not meet any further plane of $\mathcal{S}$. A similar argument shows that the lines $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$ do not meet any further plane of the regular 2-spread $\mathcal{S}$. □

6.3 Transversal lines of exterior splashes with common carriers

As exterior splashes are equivalent to covers of the circle geometry $CG(3,q)$, there are $q-1$ disjoint exterior splashes on ℓ_∞ with common carriers E_1, E_2. We show that in $\mathrm{PG}(6,q)$, the covers of these disjoint exterior splashes have common transversals.

Theorem 6.6 *Let $\mathbb{S}_0, \ldots, \mathbb{S}_{q-1}$ be $q-1$ disjoint exterior splashes on ℓ_∞ with common carriers E_1, E_2, and let exterior splash $\mathbb{S}_j$ have covers $\mathbb{C}_j$, $\mathbb{T}_j$. Then the covers $\mathbb{C}_0, \ldots, \mathbb{C}_{q-1}$ have common transversal lines $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$, and the covers $\mathbb{T}_0, \ldots, \mathbb{T}_{q-1}$ have common transversal lines $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$.*

Proof By Theorem 2.4, we can without loss of generality prove this for the order-q-subplane $\mathcal{B}$ coordinatised in Section 3. Let $\mathcal{K} = \{k \in \mathbb{F}'_{q^3} : k^{q^2+q+1} = 1\} = \{k = \tau^{i(q-1)} : 0 \leq i < q^2+q+1\}$. Recall that $\mathcal{B}$ has carriers $E_1 = (1,0,0)$, $E_2 = (0,1,0)$, and exterior splash $\mathbb{S}_0 = \{S_{k,0} = (k,1,0) : k \in \mathcal{K}\}$. Let $\mathcal{K}_j = \tau^j\mathcal{K}$, for $j = 0 \ldots, q-2$, be the $q-1$ cosets of $\mathcal{K}$ in $\mathbb{F}'_{q^3}$. Let $\mathbb{S}_j = \{S_{k,j} = (k,1,0) : k \in \mathcal{K}_j\}$, $0 \leq j \leq q-2$. Consider the homography ξ acting on ℓ_∞ that maps the point $(x,y,0)$ to $(\tau x, y, 0)$. Then ξ fixes E_1, E_2, maps $\mathbb{S}_j$ to $\mathbb{S}_{j+1}$ $(0 \leq j \leq q-3)$, and maps $\mathbb{S}_{q-2}$ to $\mathbb{S}_0$. Hence $\mathbb{S}_0, \ldots, \mathbb{S}_{q-1}$ are the $q-1$ disjoint exterior splashes on ℓ_∞ with carriers $(1,0,0)$ and $(0,1,0)$.

In $\Sigma_\infty \cong \mathrm{PG}(5,q)$, we have planes $[S_{k,j}] = \{([kx],[x]) : x \in \mathbb{F}'_{q^3}\} \in \mathbb{S}$, and define planes $[T_{k,j}] = \{([kx],[x^q]) : x \in \mathbb{F}'_{q^3}\}$, and $[C_{k,j}] = \{([kx],[x^{q^2}]) : x \in \mathbb{F}'_{q^3}\}$, for $k \in \mathcal{K}_j$. So $\mathbb{S}_j = \{[S_{k,j}], k \in \mathcal{K}_j\}$, and define $\mathbb{T}_j = \{[T_{k,j}], k \in \mathcal{K}_j\}$ and $\mathbb{C}_j = \{[C_{k,j}], k \in \mathcal{K}_j\}$. Note that $\mathbb{T}_0$, $\mathbb{C}_0$ are the covers of the exterior splash $\mathbb{S}_0$ of $\mathcal{B}$. Now consider the map θ_τ of $\mathrm{PG}(5,q)$ acting on Σ_∞ defined in Section 2.2, it maps $\mathbb{S}_j$ to $\mathbb{S}_{j+1}$; $\mathbb{T}_j$ to $\mathbb{T}_{j+1}$; and $\mathbb{C}_j$ to $\mathbb{C}_{j+1}$. Hence $\mathbb{T}_j$ and $\mathbb{C}_j$ are covers for $\mathbb{S}_j$. By Theorem 6.3, the transversal line of $\mathbb{T}_0$ is $g_{\mathbb{T}} = \langle A_1, A_2^{q^2}\rangle$. Using (1), we see that the homography θ_τ fixes $g_{\mathbb{T}}$, and so $g_{\mathbb{T}}$ is a transversal for all $\mathbb{T}_j$. So $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$ are transversal lines of $\mathbb{T}_j$ for each $j = 0, \ldots, q-2$. Similarly, $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$ are transversal lines of $\mathbb{C}_j$ for each $j = 0, \ldots, q-2$. □

Remark 6.7 We can interpret this result using the terminology of [13]. We can partition the planes of a regular 2-spread into $q-1$ disjoint hyper-reguli with common carriers. Each hyper-reguli has two replacement hyper-reguli, which correspond to our conic and tangent covers. If we replace all $q-1$ hyper-reguli of $\mathcal{S}$ with hyper-reguli of the *same type* (that is, all belonging to $\mathbb{C}$, or all belonging to $\mathbb{T}$), then the resulting 2-spread has transversals either $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$ or $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$, and so is regular. Hence the resulting André plane is Desarguesian. If we replace all the hyper-reguli of $\mathcal{S}$ with a combination of hyper-reguli from each type, then the resulting 2-spread is not regular, and so the resulting André plane is non-Desarguesian.

7 A characterisation of the sublines of an exterior splash

Let π be an exterior order-q-subplane of $\mathrm{PG}(2,q^3)$ with exterior splash $\mathbb{S}$ on ℓ_∞. The order-q-sublines of $\mathbb{S}$ are studied in [17], and are characterised with respect to geometric

objects of π in [6]. In this section we characterise the order-q-sublines of $\mathbb{S}$ with respect to the covers of $\mathbb{S}$ and their transversal lines.

There are $2(q^2+q+1)$ order-q-sublines in an exterior splash which lie in two families of size q^2+q+1. By [6, Theorem 5.2], the two families are related to the associated exterior order-q-subplane π as follows. If A is a point of π, then the pencil of $q+1$ lines of π through A meets ℓ_∞ in an order-q-subline of $\mathbb{S}$, called a *π-pencil-subline* of $\mathbb{S}$. If Γ is a (π,ℓ_∞)-special-dual conic of π, then the lines of Γ meet ℓ_∞ in an order-q-subline of $\mathbb{S}$, called a *π-dual-conic-subline* of $\mathbb{S}$. Note that we need to define these two classes with respect to an associated order-q-subplane π, since in [6, Theorem 5.3], we show that the two families of order-q-sublines of $\mathbb{S}$ can act as either class by considering different associated order-q-subplanes.

We now consider the interaction in $\mathrm{PG}(6,q)$ of these two families of order-q-sublines with the two covers of $\mathbb{S}$. We show in Theorem 7.1 that each family meets planes from one cover in lines, and planes from the other cover in conics. Further, Theorem 7.2 shows that the converse is true, and so we have a characterisation of the order-q-sublines of $\mathbb{S}$. Moreover, Theorem 7.3 shows that the conics concerned in each case are special.

Suppose $\mathcal{R}$ is a 2-regulus in $\mathrm{PG}(5,q)$, and consider a plane α that meets $\mathcal{R}$ in a set of $q+1$ points. Then an easy counting argument shows that these points form either a line or a conic in α. We abbreviate this to '$\mathcal{R}$ meets α in a line or a conic'.

Theorem 7.1 *Let π be an exterior order-q-subplane with exterior splash $\mathbb{S}$, conic cover $\mathbb{C}$, and tangent cover $\mathbb{T}$.*

1. *A π-pencil-subline of $\mathbb{S}$ corresponds in $\mathrm{PG}(6,q)$ to a 2-regulus that meets each plane of $\mathbb{T}$ in a distinct line, and meets each plane of $\mathbb{C}$ in a conic.*
2. *A π-dual-conic-subline of $\mathbb{S}$ corresponds in $\mathrm{PG}(6,q)$ to a 2-regulus that meets each plane of $\mathbb{T}$ in a conic, and meets each plane of $\mathbb{C}$ in a distinct line.*

Proof Let P be a point in the exterior order-q-subplane π, and let d be the corresponding π-pencil-subline of $\mathbb{S}$. By Theorem 2.3, in $\mathrm{PG}(6,q)$, $[d]$ is a 2-regulus contained in $\mathbb{S}$. Consider the tangent plane $\mathcal{T}_P$ to $[\pi]$ at P. By Theorem 4.2, the lines of $\mathcal{T}_P$ through P meet Σ_∞ in points that lie in distinct planes of the 2-regulus $[d]$. Hence $\mathcal{T}_P \cap \Sigma_\infty$ is a ruling line of the 2-regulus $[d]$. By Theorem 5.3, this ruling line $\mathcal{T}_P \cap \Sigma_\infty$ lies in a tangent cover plane. The homography Θ of Lemma 5.2 fixes the planes of $[b]$ and is transitive on the cover planes of $\mathbb{T}$. Hence each ruling line of $[b]$ meets a unique cover plane of $\mathbb{T}$.

A straightforward geometric argument shows that planes of $\mathbb{T},\mathbb{C}$ meet a 2-regulus of $\mathbb{S}$ in a line or a conic. Hence a conic cover plane meets the 2-regulus $[d]$ in a conic. As there are q^2+q+1 π-pencil-sublines of $\mathbb{S}$, every line in a plane of $\mathbb{T}$ is a ruling line for some 2-regulus

corresponding to a π-pencil-subline. Hence if $[d']$ is a 2-regulus of $\mathbb{S}$ corresponding to a π-dual-conic-subline, then planes of $\mathbb{T}$ meet $[d']$ in conics, and so planes of $\mathbb{C}$ meet $[d']$ in ruling lines of $[d']$. Moreover, applying the homography of Lemma 5.2 shows that each ruling line of $[d']$ lies in a unique conic cover plane. $\square$

By Theorem 2.3, there is a one to one correspondence between the order-q-sublines of $\mathbb{S}$ in $\mathrm{PG}(2, q^3)$, and the 2-reguli contained in $\mathbb{S}$ in $\mathrm{PG}(6, q)$. Hence the converse of Theorem 7.1 is also true, and so we have a characterisation of order-q-sublines of $\mathbb{S}$ relating to the cover planes of the associated order-q-subplane.

Theorem 7.2 *Let π be an exterior order-q-subplane with exterior splash $\mathbb{S}$, conic cover $\mathbb{C}$, and tangent cover $\mathbb{T}$.*

1. *A π-pencil-subline of $\mathbb{S}$ corresponds to a 2-regulus that meets each plane of $\mathbb{T}$ in a distinct line. Conversely, a 2-regulus contained in $\mathbb{S}$ that meets some plane of $\mathbb{T}$ in a line corresponds to a π-pencil-subline of $\mathbb{S}$.*

2. *A π-pencil-subline of $\mathbb{S}$ corresponds to a 2-regulus that meets each plane of $\mathbb{C}$ in a conic. Conversely, a 2-regulus contained in $\mathbb{S}$ that meets some plane of $\mathbb{C}$ in a conic corresponds to a π-pencil-subline of $\mathbb{S}$.*

3. *A π-dual-conic-subline of $\mathbb{S}$ corresponds to a 2-regulus that meets each plane of $\mathbb{T}$ in a conic. Conversely, a 2-regulus contained in $\mathbb{S}$ that meets some plane of $\mathbb{T}$ in a conic corresponds to a π-dual-conic-subline of $\mathbb{S}$.*

4. *A π-dual-conic-subline of $\mathbb{S}$ corresponds to a 2-regulus that meets each plane of $\mathbb{C}$ in a distinct line. Conversely, a 2-regulus contained in $\mathbb{S}$ that meets some plane of $\mathbb{C}$ in a line corresponds to a π-dual-conic-subline of $\mathbb{S}$.*

In fact, we can give a stronger characterisation of the order-q-sublines of $\mathbb{S}$, namely that the conics of Theorem 7.1 are special with respect to the associated cover. In order to prove that the conics are special, we need to introduce coordinates, and the proof is calculation intensive.

Theorem 7.3 *Let π be an exterior order-q-subplane with exterior splash $\mathbb{S}$, conic cover $\mathbb{C}$, and tangent cover $\mathbb{T}$.*

1. *A 2-regulus of $\mathbb{S}$ corresponding to a π-pencil-subline of $\mathbb{S}$ meets each plane of $\mathbb{C}$ in a $\mathbb{C}$-special conic.*

2. *A 2-regulus of $\mathbb{S}$ corresponding to a π-dual-conic-subline of $\mathbb{S}$ meets each plane of $\mathbb{T}$ in a $\mathbb{T}$-special conic.*

Proof By Theorem 2.4, we can without loss of generality prove this for the exterior order-q-subplane $\mathcal{B}$ coordinatised in Section 3. We start with the order-q-subplane $\pi_0 = \mathrm{PG}(2,q)$ and the line $\ell = [-\tau\tau^q, \tau+\tau^q, -1]$ which is exterior to π_0. Note that using the notation for p_0, p_1, p_2 given in Theorem 6.3, we have $\ell = [p_0^{q^2}, p_1^{q^2}, p_2^{q^2}]$. A line of π_0 has coordinates $[l,m,n]$ for $l,m,n \in \mathbb{F}_q$, and meets ℓ in the point $W'_{l,m,n} = (-n(\tau+\tau^q)-m,\ l-n\tau\tau^q,\ m\tau\tau^q + l(\tau+\tau^q))$. We apply the homography σ of Section 3 with matrix K to map π_0, ℓ to $\mathcal{B}$, ℓ_∞, respectively. The point $W'_{l,m,n}$ of ℓ maps to the point $W_{l,m,n} = KW'_{l,m,n} = (l+m\tau+n\tau^2,\ l+m\tau^q+n\tau^{2q},\ 0)$ of ℓ_∞. Writing $\varepsilon = \varepsilon_{l,m,n} = l+m\tau+n\tau^2$, we have $W_\varepsilon = W_{l,m,n} = (\varepsilon, \varepsilon^q, 0) \equiv (\varepsilon^{1-q}, 1, 0)$. Using the notation from Lemma 5.1, this is the point $S_{\varepsilon^{1-q}} \in \ell_\infty$. In $\mathrm{PG}(6,q)$, W_ε corresponds to the spread plane $[W_\varepsilon] = [W_{l,m,n}] = \{([\varepsilon x], [\varepsilon^q x], 0) \equiv ([\varepsilon^{1-q}x], [x], 0) : x \in \mathbb{F}'_{q^3}\} = [S_{\varepsilon^{1-q}}]$.

Fix a point $P = (a,b,c)$ of π_0, so $a,b,c, \in \mathbb{F}_q$, not all zero. Let $\mathcal{L} = \{(l,m,n) : l,m,n \in \mathbb{F}_q$ not all zero, and $la+mb+nc = 0\}$. The $q+1$ lines of π_0 through P have coordinates $[l,m,n] \in \mathcal{L}$. These $q+1$ lines meet the exterior line ℓ of π_0 in a π_0-pencil-subline which, under the collineation σ, maps to a $\mathcal{B}$-pencil-subline d of ℓ_∞. By Theorem 2.3, in $\mathrm{PG}(6,q)$, d corresponds to the 2-regulus $[d]$ which we denote by $\mathcal{R}$, so $\mathcal{R} = [d] = \{[W_\varepsilon] = [S_{\varepsilon^{1-q}}] : \varepsilon \in \mathcal{W}\}$, where $\mathcal{W} = \{\varepsilon = \varepsilon_{l,m,n} = l+m\tau+n\tau^2 : (l,m,n) \in \mathcal{L}\}$. For each $\alpha \in \mathbb{F}'_{q^3}$, consider the set of points $t_\alpha = \{([\varepsilon\alpha], [\varepsilon^q\alpha], 0) : \varepsilon \in \mathcal{W}\}$. As $\mathcal{W}$ is closed under addition, t_α is a line of $\Sigma_\infty \cong \mathrm{PG}(5,q)$, further t_α meets every plane in $\mathcal{R}$. Hence t_α is a ruling line of the 2-regulus $\mathcal{R}$.

By Theorem 7.2(2), the 2-regulus $\mathcal{R}$ meets a cover plane of the conic cover $\mathbb{C}$ in a conic $\mathcal{C}_k = [C_k] \cap \mathcal{R}$ for $k \in \mathcal{K}$. To show that the conic $\mathcal{C}_k$ is $\mathbb{C}$-special, we need to extend it to $\mathrm{PG}(5,q^3)$, and show that it meets the three transversal lines of $\mathbb{C}$. To do this, we extend the 2-regulus $\mathcal{R}$ of $\Sigma_\infty \cong \mathrm{PG}(5,q)$ to a 2-regulus $\mathcal{R}^*$ of $\mathrm{PG}(5,q^3)$, so $\mathcal{C}_k^* = [C_k]^* \cap \mathcal{R}^*$. We then use coordinates to show that one of the planes of $\mathcal{R}^*$ contains the transversal line $g_{\mathbb{C}}^{q^2}$ of $\mathbb{C}$, and then deduce that $\mathcal{C}_k^*$ meets $g_{\mathbb{C}}^{q^2}$.

To extend $\mathcal{R}$ to a 2-regulus $\mathcal{R}^*$ of $\mathrm{PG}(5,q^3)$, we find four lines in $\mathrm{PG}(5,q^3)$ that meet each extended plane of $\mathcal{R}$. As a 2-regulus is uniquely determined by four ruling lines in general position, we can use these four lines to define the 2-regulus $\mathcal{R}^*$. The transversal line $g_{\mathbb{S}}$ of the regular 2-spread $\mathcal{S}$ can be used as one of our ruling lines, for the other three ruling lines, we use the extended lines t_1^*, $t_\tau^*, t_{\tau^2}^*$, which each meet every plane of $\mathcal{R}$. So $\mathcal{R}^*$ is the 2-regulus of $\mathrm{PG}(5,q^3)$ determined by the four ruling lines t_1^*, $t_\tau^*, t_{\tau^2}^*, g_{\mathbb{S}}$ (which are in general position), and further $\mathcal{R}^* \cap \Sigma_\infty = \mathcal{R}$.

We now exhibit a plane γ of $\mathcal{R}^*$ that contains the transversal line $g_{\mathbb{C}}^{q^2}$ of the conic cover $\mathbb{C}$. Extend the set $\mathcal{L}$ to $\mathcal{L}^* = \{(l,m,n) : l,m,n \in \mathbb{F}_{q^3}$ not all zero, and $la+mb+nc = 0\}$. We use the matrix M_τ defined in Section 2.2, and write $M = M_\tau$. The ruling line $t_{\tau^i}^*$, $i = 0,1,2$, has points $P_{\tau^i,l,m,n}$ with $(l,m,n) \in \mathcal{L}^*$, where $P_{\tau^i,l,m,n} = l(M^i[1], M^i[1], 0) + m(M^i[\tau], M^i[\tau^q], 0) + n(M^i[\tau^2], M^i[\tau^{2q}], 0)$. Recall that the order-$q$-subline d corresponds

to the fixed point $P = (a, b, c) \in \pi_0$. Consider the following $(l, m, n) \in Ł^*$,

$$l = c\tau - b\tau^2, \quad m = a\tau^2 - c, \quad n = b - a\tau. \tag{9}$$

Note that for these l, m, n we have

$$l + m\tau + n\tau^2 = 0. \tag{10}$$

For l, m, n as in (9), consider the plane γ spanned by the three points $P_{1,l,m,n} \in t_1^*$, $P_{\tau,l,m,n} \in t_\tau^*$, $P_{\tau^2,l,m,n} \in t_{\tau^2}^*$. We first show that γ is a plane of the 2-regulus $\mathcal{R}^*$ by showing that the fourth ruling line $g_{\mathbb{S}}$ of $\mathcal{R}^*$ also meets γ. By Theorem 6.3, $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$, we show that $g_{\mathbb{S}}$ meets γ by showing that the point A_2 lies in γ. With l, m, n given by (9), consider the following point $F = p_0 P_{1,l,m,n} + p_1 P_{\tau,l,m,n} + p_2 P_{\tau^2,l,m,n}$ of γ. To simplify the notation, we use the the point $A = (p_0, p_1, p_2)^t$, and matrix $U_0 = p_0 I + p_1 M + p_1 M^2$ defined in Section 2.2, and note that $U_0[\alpha] = \alpha A$. We have

$$\begin{aligned} F &= (lU_0[1] + mU_0[\tau] + nU_0[\tau^2],\ lU_0[1] + mU_0[\tau^q] + nU_0[\tau^{2q}],\ 0) \\ &= (lA + m\tau A + n\tau^2 A,\ lA + m\tau^q A + n\tau^{2q} A,\ 0). \end{aligned}$$

By (10), $F \equiv ([0], A, 0) = A_2$, and by Lemma 2.1, $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$, so $F \in g_{\mathbb{S}} \cap \gamma$. That is, the four ruling lines $t_1^*, t_\tau^*, t_{\tau^2}^*, g_{\mathbb{S}}$ of the 2-regulus $\mathcal{R}^*$ all meet the plane γ, and so γ is a plane of $\mathcal{R}^*$.

We now show that the transversal line $g_{\mathbb{C}}^{q^2}$ of $\mathbb{C}$ lies in the plane γ of $\mathcal{R}^*$. Let $G = p_0^{q^2} P_{1,l,m,n} + p_1^{q^2} P_{\tau,l,m,n} + p_2^{q^2} P_{\tau^2,l,m,n}$, and note that $G \in \gamma$. We use the matrix $U_2 = p_0^{q^2} I + p_1^{q^2} M + p_1^{q^2} M^2$ defined in Section 2.2, and note that $U_2[\alpha] = \alpha^{q^2} A^{q^2}$, so we have

$$\begin{aligned} G &= (lU_2[1] + mU_2[\tau] + n^2 U_2[\tau^2],\ lU_2[1] + mU_2[\tau^q] + nU_2[\tau^{2q}],\ 0) \\ &= (lA^{q^2} + m\tau^{q^2} A^{q^2} + n\tau^{2q^2} A^{q^2},\ lA^{q^2} + m\tau A^{q^2} + n\tau^2 A^{q^2},\ 0). \end{aligned}$$

By (10), $G \equiv (A^{q^2}, [0], 0) = A_1^{q^2}$, so γ contains the points $G = A_1^{q^2}$ and $F = A_2$. Hence by Theorem 6.3, γ contains the transversal line $g_{\mathbb{C}}^{q^2} = \langle A_1^{q^2}, A_2 \rangle$ of $\mathbb{C}$.

We showed above that the 2-regulus $[d] = \mathcal{R}$ meets a cover plane $[C_i]$ of $\mathbb{C}$ in a conic $\mathcal{C}_i$. We want to show that $\mathcal{C}_i$ is a $\mathbb{C}$-special conic, that is, we want to show that in $\mathrm{PG}(6, q^3)$, the extended conic $\mathcal{C}_i^* = [C_i]^* \cap \mathcal{R}^*$ contains the three points $g_{\mathbb{C}} \cap [C_i]^*, g_{\mathbb{C}}^q \cap [C_i]^*, g_{\mathbb{C}}^{q^2} \cap [C_i]^*$. We have shown that the transversal line $g_{\mathbb{C}}^{q^2}$ of $\mathbb{C}$ lies in a plane γ of $\mathcal{R}^*$. As the extended cover plane $[C_i]^*$ meets the transversal line $g_{\mathbb{C}}^{q^2}$ in a unique point denoted P_i, we have $P_i = [C_i]^* \cap g_{\mathbb{C}}^{q^2} = [C_i]^* \cap \gamma \in [C_i]^* \cap \mathcal{R}^* = \mathcal{C}_i^*$. Hence $\mathcal{C}_i^*$ contains the point $g_{\mathbb{C}}^{q^2} \cap [C_i]^*$, and hence it also contains the conjugate points $g_{\mathbb{C}}^q \cap [C_i]^*, g_{\mathbb{C}} \cap [C_i]^*$. That is, the conic $\mathcal{C}_i = [C_i] \cap \mathcal{R}$ is a $\mathbb{C}$-special conic, completing the proof of part 1.

We now tackle part 2. Let β be the collineation defined in (7) acting on $\Sigma_\infty^* \cong \mathrm{PG}(5, q^3)$. Consider the line t_1 defined above which lies in the tangent cover plane $[T_1]$. By the proof

of Lemma 5.1, β maps the cover plane $[T_1] \in \mathbb{T}$ to the cover plane $[C_1] \in \mathbb{C}$. Hence the line $u_1 = \beta(t_1)$ lies in the cover plane $[C_1]$ of the conic cover $\mathbb{C}$. As u_1 meets $q+1$ planes of $\mathbb{S}$, these planes form a 2-regulus of $\mathbb{S}$ which we denote $\mathcal{R}_1$. Further, by Lemma 5.2, all the conic cover planes meet $\mathcal{R}_1$ in ruling lines of $\mathcal{R}_1$. By Theorem 2.3, the 2-regulus $\mathcal{R}_1$ corresponds to an order-q-subline d' of ℓ_∞ in $\mathrm{PG}(2,q^3)$. If d' were a $\mathcal{B}$-pencil-subline of $\mathbb{S}$, then by Theorem 7.1(1), the tangent cover planes would meet the 2-regulus $[d'] = \mathcal{R}_1$ in ruling lines of $\mathcal{R}_1$. However we have just shown that the conic cover planes meet $[d'] = \mathcal{R}_1$ in ruling lines of $\mathcal{R}_1$. Hence d' is a $\mathcal{B}$-dual-conic-subline of $\mathbb{S}$ in $\mathrm{PG}(2,q^3)$, and so we are in case 2.

As before, we want to extend the 2-regulus $\mathcal{R}_1$ of Σ_∞ to a 2-regulus $\mathcal{R}_1^*$ of $\Sigma_\infty^* \cong \mathrm{PG}(5,q^3)$. We need to determine coordinates of four ruling lines of the 2-regulus $\mathcal{R}_1$. Let $\varepsilon \in \mathcal{W}$, then the point $([\varepsilon],[\varepsilon^{q^2}],0)$ lies in u_1. Further, it lies in the spread element $[S_{\varepsilon^{1-q^2}}]$. Hence the 2-regulus $\mathcal{R}_1$ consists of the planes $\{[S_{\varepsilon^{1-q^2}}] : \varepsilon \in \mathcal{W}\}$. As $[S_{\varepsilon^{1-q^2}}]$ has elements $([\varepsilon^{1-q^2}x],[x],0) \equiv ([\varepsilon x],[\varepsilon^{q^2}x],0)$, the ruling lines of the 2-regulus $\mathcal{R}_1$ are given by $v_\alpha = \{([\alpha\varepsilon],[\alpha\varepsilon^{q^2}],0) : \varepsilon \in \mathcal{W}\}$, for $\alpha \in \mathbb{F}'_{q^3}$ (note that $u_1 = v_1$). We extend the 2-regulus $\mathcal{R}_1$ to the 2-regulus $\mathcal{R}_1^*$ uniquely determined by the four ruling lines $v_1^* = u_1^*, v_\tau^*, v_{\tau^2}^*, g_{\mathbb{S}}$, where $v_{\tau^i}^* = \{Q_{\tau^i,l,m,n} : (l,m,n) \in \mathcal{L}^*\}$, and $Q_{\tau^i,l,m,n} = l(M^i[1],M^i[1],0) + m(M^i[\tau],M^i[\tau^{q^2}],0) + n(M^i[\tau^2],M^i[\tau^{2q^2}],0)$, $i = 0,1,2$. For l,m,n as in (9), consider the plane $\delta = \langle Q_{1,l,m,n}, Q_{\tau,l,m,n}, Q_{\tau^2,l,m,n}\rangle$. Clearly δ meets the ruling lines $v_1^*, v_\tau^*, v_{\tau^2}^*$ of $\mathcal{R}_1^*$. We show that δ lies in the 2-regulus $\mathcal{R}_1^*$ by showing that the fourth ruling line $g_{\mathbb{S}}$ of $\mathcal{R}_1^*$ also meets δ. Consider the point $H = p_0Q_{1,l,m,n} + p_1Q_{\tau,l,m,n} + p_2Q_{\tau^2,l,m,n}$ of δ, then

$$\begin{aligned} H &= (lU_0[1] + mU_0[\tau] + nU_0[\tau^2],\ lU_0[1] + mU_0[\tau^{q^2}] + nU_0[\tau^{2q^2}],\ 0) \\ &= ((l + m\tau + n\tau^2)A,\ (l + m\tau^{q^2} + n\tau^{2q^2})A,\ 0). \end{aligned}$$

By (10), $H \equiv ([0],A,0) = A_2$, hence $g_{\mathbb{S}} = \langle A_1, A_2\rangle$ meets δ in the point $H = A_2$, and so the plane δ lies in the 2-regulus $\mathcal{R}_1^*$.

We now show that the transversal line $g_{\mathbb{T}}^q$ of the tangent cover $\mathbb{T}$ lies in this plane δ. Let $J = p_0^qQ_{1,l,m,n} + p_1^qQ_{\tau,l,m,n} + p_2^qQ_{\tau^2,l,m,n}$, and note that $J \in \delta$. Now

$$\begin{aligned} J &= (lU_1[1] + mU_1[\tau] + nU_1[\tau^2],\ lU_1[1] + mU_1[\tau^{q^2}] + nU_1[\tau^{2q^2}],\ 0) \\ &= ((l + m\tau^q + n\tau^{2q})A^q,\ (l + m\tau + n\tau^2)A^q,\ 0). \end{aligned}$$

By (10), $J \equiv (A^q,[0],0) = A_1^q$, thus δ contains the points $H = A_2$, $J = A_1^q$, and so by Theorem 6.3, δ contains the transversal line $g_{\mathbb{T}}^q = \langle A_2, A_1^q\rangle$ of $\mathbb{T}$. Similar to the argument for part 1 above, a plane of $\mathbb{T}$ meets the 2-regulus $[d'] = \mathcal{R}_1$ in a conic, which is $\mathbb{T}$-special conic as $g_{\mathbb{T}}^q \subset \delta \subset \mathcal{R}_1^*$. That is, a $\mathcal{B}$-dual-conic-subline of ℓ_∞ corresponds to a 2-regulus of Σ_∞ that meets tangent cover planes in $\mathbb{T}$-special conics, so part 2 is proved. $\square$

8 Conclusion

An investigation into the interaction between an exterior order-q-subplane π of $\mathrm{PG}(2,q^3)$, and its exterior splash on ℓ_∞ began in [6]. The main focus of [6] was to show that exterior splashes are projectively equivalent to scattered linear sets of rank 3, covers of circle geometries, Sherk sets of size q^2+q+1. Further, we investigated the geometric relationship between the order-q-sublines of $\mathbb{S}$ and the points of π. The current article focusses on using the Bruck-Bose representation in $\mathrm{PG}(6,q)$ to continue the study of exterior splashes, in particular their interplay with order-q-subplanes. The notion of special conics and special twisted cubics is closely tied with this interplay. This theme is continued in [7] where the authors show that (π,ℓ_∞)-special conics of π in $\mathrm{PG}(2,q^3)$ correspond to $\mathbb{C}$-special conics in $\mathrm{PG}(6,q)$.

References

[1] J. André. Über nicht-Desarguessche Ebenen mit transitiver Translationgruppe. *Math. Z.*, **60** (1954) 156–186.

[2] S.G. Barwick and W.-A. Jackson. Sublines and subplanes of $\mathrm{PG}(2,q^3)$ in the Bruck-Bose representation in $\mathrm{PG}(6,q)$. *Finite Fields Appl.,* **18** (2012) 93–107.

[3] S.G. Barwick and W.-A. Jackson. A characterisation of tangent subplanes of $\mathrm{PG}(2,q^3)$. *Designs Codes, Cryptogr.* **71** (2014) 541–545.

[4] S.G. Barwick and W.-A. Jackson. An investigation of the tangent splash of a subplane of $\mathrm{PG}(2,q^3)$. *Designs Codes, Cryptogr.* 2014, (DOI) 10.1007/s10623-014-9971-3.

[5] S.G. Barwick and W.-A. Jackson. The tangent splash in $\mathrm{PG}(6,q)$. http://arxiv.org/abs/1305.6674.

[6] S.G. Barwick and W.-A. Jackson. Exterior splashes and linear sets of rank 3. http://arxiv.org/abs/1404.1641.

[7] S.G. Barwick and W.-A. Jackson. The exterior splash in $\mathrm{PG}(6,q)$: Special conics. http://arxiv.org/abs/

[8] R.H. Bruck. Construction problems of finite projective planes. *Conference on Combinatorial Mathematics and its Applications*, University of North Carolina Press, (1969) 426–514.

[9] R.H. Bruck. Circle geometry in higher dimensions. *A Survey of Combinatorial Theory*, eds. J. N. Srivastava, et al, Amsterdam (1973) 69–77.

[10] R.H. Bruck. Circle geometry in higher dimensions II. *Geom. Dedicata* **2** (1973) 133–188.

[11] R.H. Bruck and R.C. Bose. The construction of translation planes from projective spaces. *J. Algebra*, **1** (1964) 85–102.

[12] R.H. Bruck and R.C. Bose. Linear representations of projective planes in projective spaces. *J. Algebra*, **4** (1966) 117-172.

[13] C. Culbert and G.L. Ebert. Circle Geometry and three-dimensional subregular translation planes. *Innov. Incidence Geom.*, **1** (2005) 3–18.

[14] S. Ferret and L. Storme. Results on maximal partial spreads in PG$(3, p^3)$ and on related minihypers. *Des. Codes Cryptogr.*, **29** (2003) 105–122.

[15] J.W.P. Hirschfeld. *Finite Projective Spaces of Three Dimensions.* Oxford University Press, 1985.

[16] J.W.P. Hirschfeld and J.A. Thas. *General Galois Geometries.* Oxford University Press, 1991.

[17] M. Lavrauw and G. Van de Voorde. On linear sets on a projective line. *Des. Codes Cryptogr.*, **56** (2010) 89–104.

[18] M. Lavrauw and C. Zanella. Subgeometries and linear sets on a projective line http://arxiv.org/abs/1403.5754.

[19] T.G. Ostrom. Hyper-reguli. *J. Geom.*, **48** (1993) 157–166.

[20] R. Pomareda. Hyper-reguli in projective space of dimension 5, Mostly Finite Geometries (1996), *Lecture Notes in Pure and Appl. Math.*, **190** (1997) 379–381.